

BIBLIOGRAPHY

- Agnieszka Zak-Golab, Barbara Zahorska-Markiewicz, Jozef Langfort, Michal Holecki, Piotr Kocelak, Katarzyna Mizia-Stec, Magdalena Olszanecka-Glinianowicz and Jerzy Chudek. (2010). The influence of weight loss on anaerobic threshold in obese women. *Journal of Sports Science and Medicine*, 9, 564 – 571.
- Anderson, O. (2000) You May (Mistakenly) Think This Training Method Is Old Hat. *Peak Performance*, 133, P. 1-6
- Ayşe Sarsan, Füsün Ardiç, Merih Özgen, Oya Topuz, Yurdaer Sermez The effects of aerobic and resistance exercises in obese women. Department of Physical Medicine and Rehabilitation, Faculty of Medicine, Pamukkale University, Denizli, Turkey
- Aziz, A.R. Slater, G.J. Chia, M.Y.H. & Teh K.C. (2012). Effects of Ramadan fasting on training induced adaptations to a seven-week high-intensity interval exercise programme. *Science & Sports*. Volume 27, Issue 1, February 2012, Pages 31–38.
- Beashel, P. & Taylor, J. (1996) *Advanced Studies In Physical Education And Sport*. Uk: Thomas Nelson & Sons Ltd.
- Beashel, P. & Taylor, J. (1997) *The World Of Sport Examined*. Uk: Thomas Nelson & Sons Ltd.
- Bhargava, Alok (2006). "Fiber intakes and anthropometric measures are predictors of circulating hormone, triglyceride, and cholesterol concentration in the Women's Health Trial". *Journal of Nutrition* 136 (8): 2249–2254. PMID 16857849.

- Bhargava, Alok; Guthrie, J. (2002). "Unhealthy eating habits, physical exercise and macronutrient intakes are predictors of anthropometric indicators in the Women's Health Trial: Feasibility Study in Minority Populations". *British Journal of Nutrition* 88 (6): 719–728. doi:10.1079/BJN2002739. PMID 12493094.
- Bizley, K. (1994) *Examining Physical Education*. Oxford; Heinemann Educational Publishers
- Bloomer RJ, Goldfarb AH, Wideman L, McKenzie MJ, Consitt LA. (2005). Effects of acute aerobic and anaerobic exercise on blood markers of oxidative stress. *J Strength Cond Res*. 19(2):276-85.
- Bompa, T. & Cornacchia, L. (1998) *Serious Strength Training*. Usa; Human Kinetics Publishers, Inc.
- Brewer, C (2005) *Strength And Conditioning For Games Players*. Uk; Coachwise Business Solution.
- Chacon-Mikahil MP, Forti VA, Catai AM, Szrajer JS, Golfetti R, Martins LE, Lima-Filho EC, Wanderley JS, Marin Neto JA, Maciel BC, Gallo-Júnior L. (1998). Cardiorespiratory adaptations induced by aerobic training in middle-aged men: the importance of a decrease in sympathetic stimulation for the contribution of dynamic exercise tachycardia. *Braz J Med Biol Res*. 31(5):705-12.
- Chu, D. (1996) *Explosive Power And Strength*. Usa; Human Kinetics Publishers, Inc.
- Dao, H H, Frelut, M, L., Peres, G., Bourgeois, P. & Navarro. J., (2004). Effects of a multidisciplinary weight loss intervention on anaerobic and

aerobic aptitudes in severely obese adolescents. *International Journal of Obesity* 28, 870–878.

Davis, B. Et Al. (2000) *Training For Physical Fitness*. In: Davis, B. Et Al. *Physical Education And The Study Of Sport*. Spain: Harcourt Publishers, P.121-122

Davis, B. Et Al. (2000) *Physical Education And The Study Of Sport*. Uk: Harcourt Publishers Ltd.

Delavier, F. (2001) *Strength Training Anatomy*. Usa; Human Kinetics Publishers, Inc.

Douglas L. Ballor, Jean R. Harvey-Berino, Philip A. Ades, Janet Cryan, Jorge Calles-Escandon. Contrasting effects of resistance and aerobic training on body composition and metabolism after diet-induced weight loss ☆ (1996) Sims Obesity Research Center, University of Vermont, Burlington, VT, USA

Dubach P, Myers J, Dziekan G, Goebbels U, Reinhart W, Muller P, Buser P, Stulz P, Vogt P, Ratti R. Effect of high intensity exercise training on central hemodynamic responses to exercise in men with reduced left ventricular function. *J Am Coll Cardiol*. 1997;29:1591–1598.

Fletcher GF, Balady GJ, Amsterdam EA, Chaitman B, Eckel R, Fleg J, Froelicher VF, Leon AS, Piña IL, Rodney R, Simons-Morton DA, Williams MA, Bazzarre T. Exercise standards for testing and training: a statement for healthcare professionals from the American Heart Association. *Circulation*. 2001;104:1694 –1740.

Galligan, F. Et Al. (2000) *Advanced Pe For Edexcel*. Oxford; Heinemann Educational Publishers

Geliebter, A Maher, M M., Gerace, L. B Gutin, S B Heymsfield and S A Hashim Effects of strength or aerobic training on body composition, resting metabolic rate, and peak oxygen consumption in obese dieting subjects. Obesity Research Center, St Luke's-Roosevelt Hospital, Columbia University College of Physicians and Surgeons, New York, NY 10025, USA. AG658@columbia.edu Copyright © 1997 by The American Society for Clinical Nutrition, Inc

George P. Nassis, Katerina Papantakou, Katerina Skenderi, Maria Triandafillopoulou, Stavros A. Kavouras, Mary Yannakoulia, George P. Chrousos, Labros S. Sidossis. (2005). Aerobic exercise training improves insulin sensitivity without changes in body weight, body fat, adiponectin, and inflammatory markers in overweight and obese girls. *Metabolism*. Volume 54, Issue 11, Pages 1472–1479.

Gibala MJ, McGee SL. Metabolic adaptations to short-term high-intensity interval training: a little pain for a lot of gain? *Exerc Sport Sci Rev*. 2008 Apr;36(2):58-63.

Hambrecht R, Gielen S, Linke A, Fiehn E, Yu J, Walther C, Schoene N, Schuler G. Effects of exercise training on left ventricular function and peripheral resistance in patients with chronic heart failure: a randomized trial. *JAMA*. 2000;283:3095–3101.

Helgerud J, Hoydal K, Wang E, Karlsen T, Berg P, Bjerkaas M, Simonsen T, Helgesen C, Hjorth N, Bach R, Hoff J. (2007). Aerobic high-intensity intervals improve VO₂max more than moderate training. *Med Sci Sports Exerc*. 39(4):665-71.

Jebb S. and Wells J. Measuring body composition in adults and children In: Peter G. Kopelman, Ian D. Caterson, Michael J. Stock, William H. Dietz

(2005). Clinical obesity in adults and children: In Adults and Children. Blackwell Publishing. pp. 12–28.

Kavanagh T, Mertens DJ, Hamm LF, Beyene J, Kennedy J, Corey P, Shephard RJ. Prediction of long-term prognosis in 12 169 men referred for cardiac rehabilitation. *Circulation*. 2002;106:666–671.

Kopelman P., Caterson I. An overview of obesity management In: Peter G. Kopelman, Ian D. Caterson, Michael J. Stock, William H. Dietz (2005). Clinical obesity in adults and children: In Adults and Children. Blackwell Publishing. pp. 319–326. ISBN 1-4051-1672-2.

Kraemer, William J.; Kraemer, William J.¹; Vescovi, Jason D.¹; Volek, Jeff S.¹; Nindl, Bradley C.²; Newton, Robert U.³; Patton, John F.²; Dziados, Joseph E.³; French, Duncan N.¹; Häkkinen, Keijo⁴ Effects of Concurrent Resistance and Aerobic Training on Load-Bearing Performance and the Army Physical Fitness Test. *Military Medicine*, Volume 169, Number 12, December 2004 , pp. 994-999(6)

Lafortuna CL, Agosti F, Marinone PG, Marazzi N, Sartorio A. (2004). The relationship between body composition and muscle power output in men and women with obesity. *Journal of Endocrinological Investigation*. 27(9):854-861.

Lee IM, Sesso HD, Oguma Y, Paffenbarger RS Jr. Relative intensity of physical activity and risk of coronary heart disease. *Circulation*. 2003; 107:1110–1116.

Linda M. LeMura, Serge P. von Duvillard, Joseph Andreacci, Jodi M. Klebez, Sara A. Chelland, Joseph Russo. Lipid and lipoprotein profiles, cardiovascular fitness, body composition, and diet during and after resistance, aerobic and combination training in young women.

European Journal of Applied Physiology . August 2000, Volume 82,
Issue 5-6, pp 451-458

Mcardle, W. Et Al. (2000) Essentials Of Exercise Physiology. 2nd Ed.
Philadelphia: Lippincott Williams & Wilkins

Mourot L, Bouhaddi M, Tordi N, Rouillon JD, Regnard J. Short- and long-term effects of a single bout of exercise on heart rate variability: comparison between constant and interval training exercises. Eur J Appl Physiol. 2004 Aug;92(4-5):508-17. Laboratoire de Physiologie Médecine, Faculté de Médecine et de Pharmacie, Place St. Jacques, 25030 Besançon cedex, France. mourolaurent@hotmail.com

Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. N Engl J Med. 2002;346:793– 801.

National Heart, Lung, and Blood Institute (NHLBI) (1998). Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults (PDF). International Medical Publishing, Inc. ISBN 1-58808-002-1.

Okura T, Nakata Y, Lee DJ, Ohkawara K, Tanaka K. (2005). Effects of aerobic exercise and obesity phenotype on abdominal fat reduction in response to weight loss. Int J Obes (Lond). 29(10):1259-66.

Pauletto, B. (1991) Strength Training For Coaches. Usa; Human Kinetics Publishers, Inc.

Perry CG, Heigenhauser GJ, Bonen A, Spriet LL. (2008). High-intensity aerobic interval training increases fat and carbohydrate metabolic capacities in human skeletal muscle. Appl Physiol Nutr Metab. 33(6):1112-23.

- Puhl R., Henderson K., and Brownell K. Social consequences of obesity In: Peter G. Kopelman, Ian D. Caterson, Michael J. Stock, William H. Dietz (2005). *Clinical obesity in adults and children: In Adults and Children*. Blackwell Publishing. pp. 29–45. ISBN 1-4051-1672-2.
- Rognmo O, Hetland E, Helgerud J, Hoff J, Slordahl SA. High intensity aerobic interval exercise is superior to moderate intensity exercise for increasing aerobic capacity in patients with coronary artery disease. *Eur J Cardiovasc Prev Rehabil*. 2004;11:216–222.
- Ross E. Andersen, Thomas A. Wadden, Susan J. Bartlett, Babette Zemel, Tony J. Verde, Shawn C. Franckowiak (1999). Effects of Lifestyle Activity vs Structured Aerobic Exercise in Obese Women. A Randomized Trial. *JAMA*. 281(4):335-340.
- Salvadori A, Fanari P, Marzullo P, Codecasa F, Tovaglieri I, Cornacchia M, Brunani A, Luzi L, Longhini E. (2014). Short bouts of anaerobic exercise increase non-esterified fatty acids release in obesity. *Eur J Nutr*. 2014 Feb;53(1):243-9.
- Sayyed Mohammad Marandi, Neda Ghadiri Bahram Abadi, Fahimeh Esfarjani, Hosein Mojtahedi, and Gholamali Ghasemi (2013). effects of light and moderate aerobic intensity on body composition and serum lipid profile in obese/overweight women living in Isfahan. *Int J Prev Med*. Apr 2013; 4(Suppl 1): S118–S125.
- Scholich, M. (1999) *Circuit Training For All Sports*. Canada; Sport Book Publishers
- Seidell JC. Epidemiology — definition and classification of obesity In: Peter G. Kopelman, Ian D. Caterson, Michael J. Stock, William H. Dietz

(2005). Clinical obesity in adults and children: In Adults and Children. Blackwell Publishing. pp. 3–11. ISBN 1-4051-1672-2.

Shalaby MN, Saad M, Akar S, Reda MA, Shalgham A. (2012). The Role of Aerobic and Anaerobic Training Programs on CD(34+) Stem Cells and Chosen Physiological Variables. *J Hum Kinet.* 35:69-79.

Tabata I, Nishimura K, Kouzaki M, Hirai Y, Ogita F, Miyachi M, Yamamoto K. Effects of moderate-intensity endurance and high-intensity intermittent training on anaerobic capacity and VO₂max. *Med Sci Sports Exerc.* 1996 Oct;28(10):1327-30. Department of Physiology and Biomechanics, National Institute of Fitness and Sports, Kagoshima Prefecture, Japan.

Talanian JL, Galloway SD, Heigenhauser GJ, Bonen A, Spriet LL. (2007). Two weeks of high-intensity aerobic interval training increases the capacity for fat oxidation during exercise in women. *J Appl Physiol* (1985). 102(4):1439-47.

Tancred, B. (1995) Key Methods Of Sports Conditioning. *Athletics Coach*, 29 (2), P. 19

Tenke, Z. & Higgins, A. (1999) *Medicine Ball Training*. Canada; Sport Book Publishers

Tjonna AE, Stølen TO, Bye A, Volden M, Slørdahl SA, Odegård R, Skogvoll E, Wisloff U. (2009). Aerobic interval training reduces cardiovascular risk factors more than a multitreatment approach in overweight adolescents. *Clin Sci (Lond).* 116(4):317-26.

Tomlin DL, Wenger HA. The relationship between aerobic fitness and recovery from high intensity intermittent exercise. *Sports Med.* 2001;31(1):1-11. University of Victoria, British Columbia, Canada. donat@uvic.ca

Trapp EG, Chisholm DJ, Boutcher SH. Metabolic response of trained and untrained women during high-intensity intermittent cycle exercise. *Am J Physiol Regul Integr Comp Physiol.* 2007 Dec;293(6):R2370-5. Epub 2007 Sep 26. School of Medical Sciences, Faculty of Medicine, Univ. of New South Wales, Sydney 2052, Australia. e.trapp@unsw.edu.au

Trapp EG, Chisholm DJ, Freund J, Boutcher SH. The effects of high-intensity intermittent exercise training on fat loss and fasting insulin levels of young women. *Int J Obes (Lond).* 2008 Apr;32(4):684-91. doi: 10.1038/sj.ijo.0803781. Epub 2008 Jan 15. Faculty of Medicine, University of New South Wales, Sydney, New South Wales, Australia. e.trapp@unsw.edu.au

Tremblay A, Simoneau JA, Bouchard C. Impact of exercise intensity on body fatness and skeletal muscle metabolism. *Metabolism.* 1994 Jul;43(7):814-8. Physical Activity Sciences Laboratory, Laval University, Ste-Foy, Quebec, Canada.

Whyte LJ, Gill JM, Cathcart AJ. Effect of 2 weeks of sprint interval training on health-related outcomes in sedentary overweight/obese men. *Metabolism.* 2010 Oct;59(10):1421-8.

Winch, M. (2004) *Strength Training For Athletes.* Uk; Stanley L. Hunt Ltd.

Wisloff U, Loennechen JP, Currie S, Smith GL, Ellingsen O. Aerobic exercise reduces cardiomyocyte hypertrophy and increases contractility, Ca₂₊

sensitivity and SERCA-2 in rat after myocardial infarction. *Cardiovasc Res.* 2002;54:162–174.

Wisloff U, Støylen A, Loennechen JP, Bruvold M, Rognum O, Haram PM, Tjønnå AE, Helgerud J, Slordahl SA, Lee SJ, Videm V, Bye A, Smith GL, Najjar SM, Ellingsen O, Skjaerpe T. (2007). Superior cardiovascular effect of aerobic interval training versus moderate continuous training in heart failure patients: a randomized study. *Circulation.* 19;115(24):3086-94.

World Health Organization (WHO) (2000). Technical report series 894: Obesity: Preventing and managing the global epidemic. (PDF). Geneva: World Health Organization. ISBN 92-4-120894-5.