

BIBLIOGRAPHY

BIBLIOGRAPHY**BOOKS AND PERIODICALS**

- Alan Stall G. (1980) **Encyclopedia of Physical Education, Fitness, Sports Training, Environment Nutrition**, Salt Lake City, Utah: Brighten Publishing Company, p. 419.
- Bland JM and Altman DG (1986), **Statistical Methods for Assessing Agreement between two Methods of Clinical Measurement**. Lancet, pp. 307-310
- Fix, James D. et.al. (2002). **Neuroanatomy**. Lippincott Williams & Wilkins. p. 127. ISBN 0-7817-2829-0.
- Mosby's Medical (1994), **Nursing and Allied Health Dictionary**, Fourth Edition, Mosby-Year Book , p. 1285
- Siegel, Allan (2010). **Essential Neuroscience**. Lippincott Williams & Wilkins. p. 263.
- Swenson RS. (2008) **Review of Clinical and Functional Neuroscience**, Chapter 7A: Somatosensory Systems. (online version Dartmouth college).
- Todd Ellenbecker, George J Davies (2001) **Closed Kinetic Chain Exercise. A Comprehensive Guide to Multiple Joint Exercises**. Human Kinetics, 1st Edition, ISBN 978-0-7360-0170-0
- Tudor O. Bompá, (1999). **Periodization: Theory and Methodology of Training** [4th ed], Champaign, Illinois: Human Kinetics Publishers, PP. 3 – 4.
- Uppal,A.K. (2004) **Foundations of Physical Education**, Baroda: Friends Publications, P.2
- Yobu, A. (1998), **Tests and Measurements in Physical Education**, Madras: Grace Printers.

JOURNALS

- Anderson K, Behm DG (2005) "The intact of instability resistance training on balance and stability." **Sports Med**; 35(1) 45-53
- Behm DG, Anderson K, and Curnew RS (2002). "Muscle force and activation under stable and unstable conditions" **J Str Cond Res**; 16: 416-422.
- Bird M, et.al.(2001) "The long-term benefits of a multi-component exercise intervention to balance and mobility in healthy older adults". **Arch Gerontol Geriatr**; 52(2):211-6
- Blackburn JR, Morrissey MC (1998) "The relationship between open and closed kinetic chain strength of the lower limb and jumping performance." **J Orthop Sports Phys Ther**; 27(6):430-5
- Bland JM and Altman DG, (1999) "Measuring agreement method comparison studies." **Statistical Methods in Medical Research**; 8: 135-160
- Borsa P.A.et al, (1994) "Function assessment and rehabilitation of shoulder proprioception for glenohumeral instability", **J. Sports. Rehab**; 3:84-104
- Bradley PS, Olsen PD, Portas MD (2007) "The Effect of Static, Ballistic, and Proprioceptive Neuromuscular Facilitation Stretching on Vertical Jump Performance" **Journal of Strength Conditioning Research**; 21(1): 223-6.
- Caplan,N. et. al. (2009), "The Effect of Proprioceptive Neuromuscular Facilitation and Static Stretch Training on Running Mechanics" **Journal of Strength Conditioning Research**; 23(4): 1175-80.
- Cappellino F, et.al. (2012), "Neurocognitive rehabilitative approach effectiveness after anterior cruciate ligament reconstruction with patellar tendon.", **Eur J Phys Rehabil Med.**; 48(1):17-30.
- Castro CM, King AC and Brassington GS. (2001), "Telephone versus mail interventions for maintenance of physical activity in older adults". **Health Psychol.**;20(6):438-44

- Cervantes SJ, and Snyder AR. (2011), "The effectiveness of a dynamic warm-up in improving performance in college athletes"; **J Sport Rehabil.**; 20(4):487-93
- Cressey, E.M., et al. (2007). "The effects of ten weeks of lower-body unstable surface training on markers of athletic performance"; **J Str Cond Res.**; 21(2): 561-567.
- Cressman EK, and Henriques DY. (2011), "Motor adaptation and proprioceptive recalibration." **Prog Brain Res**; 191:91-9.
- Fitzgerald GK (1997), "Open versus closed kinetic chain exercise: issues in rehabilitation after anterior cruciate ligament reconstructive surgery", **Phys Ther.**; 77(12):1747-54.
- Franco NR, et.al. (2012), " Effects of proprioceptive training program on core stability and center of gravity control in sprinters.", **J Strength Cond Res.**; 26(8):2071-7.
- GarnSN, Newton RA (1988) "Kinesthetic Awareness in Subjects with Multiple Ankle Sprains," **Phys Ther**; 68: 1667-71.
- Gillespie LD, et al. (2009) "Interventions for preventing falls in older people living in the community." **Cochrane Database Syst Rev.**; (2):CD007146.
- Glencross, D. and Thornton, E. (1981) "Position Sense Following Joint Injury", **Journal of Sports Medicine and Physical Fitness**; 21(1): 23-27
- Gomes TM, et.al. (2011), "Acute effects of two different stretching methods on local muscular endurance performance." **J Strength Cond Res**; 25(3):745-52
- Graham VL, Gehlsen GM, Edwards JA. *J Athl Train.* (1993) "Electromyographic evaluation of closed and open kinetic chain knee rehabilitation exercises". **J Athl Train**; 28(1):23-30
- Guilhem G, et.al. (2010) " Neuromuscular and muscle-tendon system adaptations to isotonic and isokinetic eccentric exercise." **Ann Phys Rehabil Med.**; 53(5):319-41

- Harmer PA, Li F (2008). "Tai Chi and falls prevention in older people". **Med Sport Sci**; 52:124–34
- Hoogenboom BJ et.al. (2009), "Using rolling to develop neuromuscular control and coordination of the core and extremities of athletes." **N Am J Sports Phys Ther.**; 4(2):70-82.
- Hrysomallis C.(2007) "Relationship between balance ability, training and sports injury risk". **Sports Med**; 37(6):547–56.
- Huang H, et.al. (2006), "Recent developments in biofeedback for neuromotor rehabilitation.", **J Neuroeng Rehabil.**; 21;3:11.
- Hubbard IJ, et.al. (2009), "Task-specific training: evidence for and translation to clinical practice.", **Occup Ther Int.**;16(3-4):175-89
- Janssen KW, et.al. (2011), "Ankles back in randomized controlled trial (ABrCt): braces versus neuromuscular exercises for the secondary prevention of ankle sprains. Design of a randomised controlled trial.", **BMC Musculoskelet Disord.**; 27;12:210
- Judge LW, et.al. (2003), "Neural adaptations with sport-specific resistance training in highly skilled athletes." **J Sports Sci.**; 21(5):419-27
- Koceja DM, (2004), "Neuromuscular characteristics of endurance- and power-trained athletes.", **Res Q Exerc Sport.**; 75(1):23-30.
- Leanderson J, Eriksson E, Nilsson C, Wykman A. (1966) "Proprioception in classical ballet dancers: a prospective study of the influence of an ankle sprain on proprioception in the ankle joint". **Am J Sports Med.**; 24:370-374.).
- Lephart SM and Henry DJ (1995) "Functional Rehabilitation of the Upper and lower extremity" **Orthop. Clin. North. Am.**; 26:579-92
- Levac D, et.al. (2009) "The application of motor learning strategies within functionally based interventions for children with neuromotor conditions." **Pediatr Phys Ther**; 21(4):345-55

- Loren Seagrave (1998) cited by Jimson Lee (2009), "Loren Seagrave – Neuro Biochemics of Maximum Velocity", <http://speedendurance.com>.
- Lucas, R.C., Koslow R. (1984) "Comparative Study of Static, Dynamic, and Proprioceptive Neuromuscular Facilitation Stretching Techniques on Flexibility" **Percept Mot Skills**; 58(2), : 615-8.
- Masci I, et.al. (2010), "Neuromechanical evidence of improved neuromuscular control around knee joint in volleyball players." **Eur J Appl Physiol.**; 108(3):443-50.
- McNeal JR, and Sands WA. (2006), "Stretching for performance enhancement." **Curr Sports Med Rep.**; 5(3):141-6.
- Mörl F, et.al. (2011), "Pain relief due to physiotherapy doesn't change the motor function of the shoulder." **J Bodyw Mov Ther**; 15(3):309-18
- Myer GD, Ford KR, McLean SG, Hewett TE. (2006) "The effects of plyometric versus dynamic stabilization and balance training on lower extremity biomechanics." **The American Journal of Sports Medicine**; 34(3): 445-455.
- Niemeijer AS, et.al. (2007), "Neuromotor task training for children with developmental coordination disorder: a controlled trial." **Dev Med Child Neurol**; 49(6):406-11.
- Osternig, L.R. et. al. (1987), "Muscle activation during Proprioceptive Neuromuscular Facilitation (PNF) stretching techniques." **American Journal of physical Medicine**; 66(5): 298-307.
- Pereira MP. (2012), "Proprioceptive neuromuscular facilitation does not increase blood pressure of healthy elderly women." **Physiother Theory Pract.**; 28(5): 412-16.
- Place N, et.al. (2013), "Effects of a short proprioceptive neuromuscular facilitation stretching bout on quadriceps neuromuscular function, flexibility, and vertical jump performance.", **J Strength Cond Res.**; 27(2):463-70

- Prokopy MP, et.al.(2008) "Closed-kinetic chain upper-body training improves throwing performance of NCAA Division I softball players." **J Strength Cond Res.**; 22(6):1790-8.
- Rammelt S, et.al. (2011), "Ligamentous injuries to the ankle joint", **Z Orthop Unfall.**;149(5):e45-67
- Rowlands,A.V. V.F. Marginson and J. Lee (2003), "Chronic Flexibility Gains: Effect of Isometric Contraction Duration During Proprioceptive Neuromuscular Facilitation Stretching Techniques" **Research Quarterly Exercises Sport**; 74(1): 47-51.
- Ryan EE, et.al. (2010), "The effects of the contract-relax-antagonist-contract form of proprioceptive neuromuscular facilitation stretching on postural stability.", **J Strength Cond Res.**; 24(7):1888-94.
- Salomonczyk D, et.al. (2011), "Proprioceptive recalibration following prolonged training and increasing distortions in visuomotor adaptation." **Neuropsychologia.**; 49(11):3053-62.
- Salsabili H, et.al. (2011), "Dynamic stability training improves standing balance control in neuropathic patients with type 2 diabetes." **J Rehabil Res Dev**; 48(7):775-86
- Schoemaker MM, et.al. (2003), "Effectiveness of neuromotor task training for children with developmental coordination disorder: a pilot study." **Neural Plast**; 10(1-2):155-63
- Seto JL, Brewster CE (1994) "Treatment approaches following foot and ankle injury". **Clin Sports Med.**;13(4):695-718.
- Shanthi Johnson, Krista Leck (2010). "The effects of dietary fasting on physical balance among healthy young women" **Nutrition Journal**, 9:18
- Sherrington CS (1907). "On the proprioceptive system, especially in its reflex aspect" **Brain** 29 (4): 467-85.
- Shimura, K. and T. Kasai, (2002) "Effects of Proprioceptive Neuromuscular Facilitation on the Initiation of Voluntary Movement and Motor Evoked

- Potentials in Upper Limb Muscles” *Human Movement Science*; 21(1), 101-13.
- Shumway-Cook A, Anson D, and Haller S. (1988) "Postural sway biofeedback: its effect on reestablishing stance stability in hemiplegic patients". *Arch. Phys. Med. Rehabil.*; 69 (6): 395–400.
- Sporis G, Jukic I, Milanovic L, Vucetic V. (2010) “Reliability and factorial validity of agility tests for soccer players.” *J Strength Cond Res*; 24(3):679-86.
- Taube W, et.al. (2007), „Differential reflex adaptations following sensorimotor and strength training in young elite athletes” *Int J Sports Med.*; 28(12):999-1005
- Thornton EW, Sykes KS, Tang WK.(2004) “Health benefits of tai chi exercise: improved balance and blood pressure in middle-aged women” *Health Promot Int.*; 19(1):33–8.
- Wheaton LA, et.al. (2007), “How does the brain respond to unimodal and bimodal sensory demand in movement of the lower extremity?” *Exp Brain Res.*; 180(2):345-54.
- Willardson JM (2004). “The effectiveness of resistance exercises performed on unstable equipment”. *Strength and Conditioning Journal*; 26 (5), 70-74.
- Wong YM, and Ng G. (2010), “Resistance training alters the sensorimotor control of vasti muscles.”, *J Electromyogr Kinesiol.*; 20(1):180-4.
- Wu G. (2002) “Evaluation of the effectiveness of tai chi for improving balance and preventing falls in the older population-a review.” *J Am Geriatr Soc.*; 50(4):746–54